



(bix-LEN-vo)

ABOUT BIXLENVO

BIXLENVO is a prescription medicine used to treat HIV-1 in adults to replace their current HIV-1 medicines when their healthcare provider determines that they meet certain requirements.

BIXLENVO does not cure HIV-1 or AIDS. HIV-1 is the virus that causes AIDS.

Do not take BIXLENVO if you take dofetilide or certain other medications called strong CYP3A inducers. Ask your healthcare provider if you are not sure.

BEFORE TAKING BIXLENVO

Tell your healthcare provider all your medical conditions, including if you:

- Have hepatitis B virus (HBV).
- Are pregnant or plan to become pregnant.
- Are breastfeeding or plan to breastfeed. Talk to your healthcare provider about the risks to your baby from breastfeeding during treatment with BIXLENVO.

Tell your healthcare provider about all the medicines you take:

- Keep a list of all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements, including St. John's wort, and show it to your healthcare provider and pharmacist.
- Ask your healthcare provider or pharmacist about medicines that interact with BIXLENVO.
- Do not start a new medicine without telling your healthcare provider.

POSSIBLE SIDE EFFECTS OF BIXLENVO

The most common side effects of BIXLENVO in clinical studies were headache (4%), nausea (3%), and diarrhea (2%). These are not all the possible side effects of BIXLENVO. Call your doctor for medical advice about side effects.

IMPORTANT FACTS

This is only a brief summary of important information about BIXLENVO™ and does not replace talking to your healthcare provider about your condition and your treatment

HOW TO TAKE BIXLENVO

- Your treatment includes BIXLENVO tablets and, **for the first two days only**, another separate medicine called SUNLENCA® tablets.
- Take BIXLENVO and SUNLENCA tablets by mouth, with or without food.

Your dosing schedule will start as follows:

- On **Day 1**, take 1 BIXLENVO tablet and 2 SUNLENCA tablets.
- On **Day 2**, take 1 BIXLENVO tablet and 2 SUNLENCA tablets.
- **It is very important that you take BIXLENVO with SUNLENCA on these first two days.** Contact your healthcare provider if you have questions.

Your dosing schedule will then continue as follows:

- On **Day 3 and each day after**, take 1 BIXLENVO tablet each day.
- If you miss a dose of BIXLENVO or SUNLENCA on Day 1 or Day 2, take it as soon as possible. Do not take Day 1 and Day 2 doses of SUNLENCA on the same day.
- If you miss a dose of BIXLENVO on Day 3 and each day after, take it as soon as possible. If more than 7 days have passed since your last dose of BIXLENVO, contact your healthcare provider as your healthcare provider will need to determine how to continue your HIV-1 treatment.

GET MORE INFORMATION

- This is only a brief summary of important information about BIXLENVO. Talk to your healthcare provider or pharmacist to learn more.
- Go to [BIXLENVO.com](https://www.bixlenvo.com) or call 1-800-GILEAD-5.
- If you need help paying for your medicine, visit [BIXLENVO.com](https://www.bixlenvo.com) for program information.



BIXLENVO, the BIXLENVO Logo, SUNLENCA, GILEAD, and the GILEAD Logo are trademarks of Gilead Sciences, Inc., or its related companies. All other marks referenced herein are the property of their respective owners.

© 2026 Gilead Sciences, Inc. All rights reserved. US-BLNC-0004 08/26